

## **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07991 - Goat Anti-Acylglycerol kinase Antibody 🏹

Size: 100µg specific antibody in 200µl

# **Target Protein**

Principal Names: acylglycerol kinase, FLJ10842, MULK, multi-substrate lipid kinase, multiple substrate lipid kinase, AGK Official Symbol: AGK Accession Number(s): NP\_060708.1 Human GenelD(s): <u>55750</u> Non-Human GenelD(s): 69923 (mouse)

#### Immunogen

Peptide with sequence CDPRKREQMLTSP, from the C Terminus of the protein sequence according to NP\_060708.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

**Western blot:** Approx 50kDa band observed in Human Brain (Substantia nigra, Hippocampus) lysates (calculated MW of 47.1kDa according to NP\_060708.1). Recommended concentration: 1-3µg/ml. An additional band of unknown identity was also consistently observed at 22kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

### **Species Reactivity**

Tested: Human Expected from sequence similarity: Human, Mouse

_	250kDa 150kDa 100kDa 75kDa 50kDa
	37kDa
	25kDa 20kDa
	15kDa

EB07991 (1µg/ml) staining of Human Brain (Substantia nigra) lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.