

#### **UK Office**

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

## **US Office**

#### Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06598 - Goat Anti-ABAD / HADH2 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: HADH2, ABAD, ERAB, MHBD, HSD17B10, 17b-HSD10, hydroxyacyl-Coenzyme A dehydrogenase, type II, type 10 17b-HSD, AB-binding alcohol dehydrogenase, type 10 17beta-hydroxysteroid dehydrogenase Official Symbol: HSD17B10 Accession Number(s): NP\_004484.1; NP\_001032900.1 Human GeneID(s): <u>3028</u> Important Comments: This antibody is expected to recognise both reported isoforms.

#### Immunogen

Peptide with sequence CIRLDGAIRMQP, from the C Terminus of the protein sequence according to NP\_004484.1; NP\_001032900.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 25kDa band observed in Human Brain lysates (calculated MW of 26.9kDa according to NP\_004484). Recommended concentration: 0.3-2µg/ml. Primary incubation 1 hour at room temperature. A minor band of unknown identity was also consistently observed at appr 38kDa. This band was successfully blocked by incubation with the immunising peptide.

### **Species Reactivity**

Tested: Human Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

	250kDa 150kDa 100kDa 75kDa 50kDa
	37kDa
-	25kDa
	20kDa
	15kDa
	10kDa

EB06598 staining (0.3µg/ml) of Human Brain lysate (RIPA buffer, 30µg total protein per lane). Detected by chemiluminescence.