

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09858-B - Goat Anti-Aconitase 2 (aa541-555), Biotinylated Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** ACO2, aconitase 2, ACONM, HEL-S-284, ICRD, OCA8, OPA9, aconitase 2, mitochondrial, citrate hydro-lyase, epididymis secretory sperm binding protein

Li 284, mitochondrial aconitase

Official Symbol: ACO2

Accession Number(s): NP\_001089.1

Human GeneID(s): 50

#### **Immunogen**

Peptide with sequence CQDTYQHPPKDSSGQH., from the internal region of the protein sequence according to NP\_001089.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

**Western blot:** Approx 90kDa band observed in Human Heart lysates, in Human, Mouse and Rat Adipose lysates and in Mouse, Rat and Pig Skeletal Muscle lysates (calculated MW of 85.4kDa according to NP\_001089.1). Recommended concentration:

0.01-0.03µg/ml. See non-biotinylated parental product's datasheet for further QC data

# **Species Reactivity**

Tested: Human, Mouse, Rat, Pig

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow, Pig

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

Biotinylated EB09858 (0.01µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.