

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire **OX25 5HD**

Enquiries:

info@everestbiotech.com

Sales:

UK

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona,

CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08207-B - Goat Anti-ACVR1, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ACVR1, activin A receptor type 1, ACTRI, ACVR1A, ACVRLK2, ALK2, FOP, SKR1, TSRI, TGF-B superfamily receptor type I, activin A receptor type I, activin A receptor, type I, activin A receptor, type II-like kinase 2, activin receptor type I, activin receptor-like kinase 2, hydroxyalkyl-protein kinase, serine/threonine-protein kinase

receptor R1

Official Symbol: ACVR1

Accession Number(s): NP_001096.1

Human GeneID(s): 90

Non-Human GenelD(s): 11477 (mouse), 79558 (rat)

Important Comments: Reported variants represent identical protein: NP_001096.1,

NP_001104537.1

Immunogen

Peptide with sequence C-RKFKRRNQERLNPRD., from the internal region of the protein sequence according to NP_001096.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 60kDa band observed in Human Umbilical Cord lysates (calculated MW of 57.2kDa according to NP_001096.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

Biotinylated EB08207 (0.1μg/ml) staining of Human Umbilical Cord lysate (35μg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.