

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07612 - Goat Anti-ADH5 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ADH5, alcohol dehydrogenase 5 (class III), chi polypeptide, ADH-3, ADHX, FDH, Alcohol dehydrogenase (class III), chi polypeptide, class III alcohol dehydrogenase 5 chi subunit, formaldehyde dehydrogenase, glutathione-dependent formaldehyde dehydrogenase, GSNOR

Official Symbol: ADH5

Accession Number(s): NP_000662.3

Human GeneID(s): 128

Non-Human GenelD(s): 11532 (mouse)

Immunogen

Peptide with sequence C-KKIKVDEFVTHN, from the internal region of the protein sequence according to NP_000662.3 .

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.

Western blot: Approx 38kDa band observed in Human Testis lysates (calculated MW of 39.7kDa according to NP_000662.3). Recommended concentration: 0.5-1.5μg/ml. An anonymous customer found positive results in WB on Rat penis tissue. An anonymous customer found positive results in WB on Mouse penile tissue.

IHC: An anonymous customer found positive results in IHC on Rat penile tissue.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

250kDa 150kDa 100kDa 75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

EB07612 (0.5μg/ml) staining of Human Testis lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.