

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona,

CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08336 - Goat Anti-AKR1B10 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: AKR1B10, aldo-keto reductase family 1, member B10 (aldose reductase), AKR1B11, AKR1B12, ALDRLn, ARL-1, ARL1, HIS, HIS, MGC14103, aldo-keto reductase family 1, member B10, aldo-keto reductase family 1, member B11 (aldose reductase-like), aldose reductase-like 1N: aldose reductase-like peptide, aldose reductase-related protein, small intestine reductase

Official Symbol: AKR1B10

Accession Number(s): NP_064695.2

Human GeneID(s): 57016

Immunogen

Peptide with sequence C-QSSHLEDYPFDAE, from the C Terminus of the protein sequence according to NP_064695.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 35kDa band observed in lysates of cell lines HEK293, A549 and HepG2 (calculated MW of 36.0kDa according to NP_064695.2). In transfected HEK293 transiently expressing AKR1B10 a band of approx. 40kDa is observed. This band is not observed in the non-transfected HEK293. Recommended concentration: 0.03-0.1µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB08336 ($0.03\mu g/ml$) staining of A549 cell lysate ($35\mu g$ protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



HEK293 overexpressing AKR1B10 (RC203177) with C-terminal tag (DYKDDDDK) and probed with anti-DYKDDDDK in the left panel and with EB08336 in the right panel (mock transfection in first and last lanes), tested by Origene.