

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12279 - Goat Anti-alpha-1-antitrypsin Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SERPINA1, serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 1, A1A, A1AT, AAT, PI, PI1, PRO2275, alpha1AT, alpha-1 protease inhibitor, alpha-1-antiproteinase, alpha-1-antitrypsin, alpha-1-antitrypsin null, protease inhibitor 1 (anti-elastase), alpha-1-antitrypsin, serine (or cysteine) proteinase inhibitor, clade A, member 1, serpin A1

Official Symbol: SERPINA1

Accession Number(s): NP_000286.3

Human GeneID(s): [5265](#)

Important Comments: The immunizing peptide represents the N terminus of the mature protein. Reported variants represent identical protein: NP_001121174.1, NP_001121179.1, NP_001121172.1, NP_001121177.1, NP_001121178.1, NP_001121176.1, NP_001121175.1, NP_000286.3, NP_001121173.1, NP_001002236.1, NP_001002235.1.

Immunogen

Peptide with sequence C-QKTDTSHHDDQDHP, from the internal region (near N terminus) of the protein sequence according to NP_000286.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 55kDa band observed in Albumin-depleted Human Plasma lysates (calculated MW of 46.7kDa according to NP_000286.3). The observed molecular weight corresponds to the glycosylated form. Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB12279 (0.3 μ g/ml) staining of Albumin-depleted Human Plasma lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.