

#### **UK Office**

**Everest Biotech Ltd** 

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10633 - Goat Anti-Ascl3 (mouse) Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** Ascl3, achaete-scute complex homolog 3 (Drosophila), MGC129412, MGC129413, Sgn1, bHLHa42, ASH-3, ICRFP703B1614Q5.4, ICRFP703N2430Q5.4, achaete-scute homolog 3, bHLH transcriptional regulator Sgn-1, mASH-3, mASH3

Official Symbol: Ascl3

Accession Number(s): NP\_064435.1

Human GeneID(s): 56676

Non-Human GenelD(s): 56787 (mouse)

#### **Immunogen**

Peptide with sequence C-HLPEDYLEKRLSK, from the internal region of the protein sequence according to NP\_064435.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:8000.

**Western blot:** Approx 23kDa band observed in Mouse Liver lysates (calculated MW of 20.2kDa according to NP\_064435.1). Recommended concentration: 0.1-0.3μg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Brain (Cortex). Recommended concentration: 3.75µg/ml.


# Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB10633 (0.1μg/ml) staining of Mouse Liver lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB10633 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.