



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07654 - Goat Anti-AVPR1A Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: AVPR1A, arginine vasopressin receptor 1A, AVPR1, SCCL vasopressin subtype 1a receptor, V1-vascular vasopressin receptor AVPR1A, V1a vasopressin receptor, antidiuretic hormone receptor 1A, vascular/hepatic-type arginine vasopressin receptor

Official Symbol: AVPR1A

Accession Number(s): NP_000697.1

Human GeneID(s): [552](#)

Non-Human GeneID(s): 54140 (mouse), 25107 (rat)

Immunogen

Peptide with sequence CHPLKTLQQPARRSR, from the internal region of the protein sequence according to NP_000697.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 85kDa band observed in Human Brain (Frontal cortex) lysates and in Rat Brain lysates and in lysates of cell line NIH3T3 (calculated MW of 46.8kDa according to NP_000697.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Phalipou et al, J Biol Chem. 1997 Oct 17;272(42):26536-44; PMID: 9334232). Recommended concentration: 0.03-0.1µg/ml.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB07654 (0.03µg/ml) staining of NIH3T3 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.