

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona,

CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08786 - Goat Anti-AVPR1B (mouse) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: AAvpr1b, arginine vasopressin receptor 1B, AVPR3, antidiuretic hormone receptor 1B, arginine vasopressin receptor 3, pituitary vasopressin receptor 3, vasopressin V1B receptor, AVPR3, V3/V1b, VIBR, VPR, V3/V1b pituitary vasopressin

receptor, VPR3

Official Symbol: AVPR1B

Accession Number(s): NP_036054.1

Non-Human GenelD(s): 26361 (mouse), 29462 (rat)

Immunogen

Peptide with sequence C-KPAGSLKDLEQVD, from the internal region (near C Terminus) of the protein sequence according to NP_036054.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx. 170kDa band observed in Rat Brain lysates (calculated MW of 46.5kDa according to NP_036054.1). The observed molecular weight corresponds to findings with EB08785, a product of different design reacting to the C-terminus of the Human protein. Recommended concentration: 0.05-0.2μg/ml.

Species Reactivity

Tested: Rat

Expected from sequence similarity: Mouse, Rat



EB08785 (0.5μg/ml) staining of Human Amygdala lysate (35μg protein in RIPA buffer) in lane A and EB08786 (0.05μg/ml) staining of Rat Brain lysate (35μg protein in RIPA buffer) in lane B . Primary incubation was 1 hour.

Detected by chemiluminescence.