



Everest Biotech Ltd
Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD, United Kingdom

everestbiotech.com

sales@everestbiotech.com

support@everestbiotech.com

Tel +44 1869 238326

Fax +44 1869 238327

Research Use Only. Not for diagnostic or therapeutic use.

Storage: For long-term storage keep aliquots at -20°C. (Store no longer than 12 months at 4°C). Minimize freezing and thawing.

EB05241 - Goat Anti-B56 beta isoform Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: protein phosphatase 2, regulatory subunit B (B56), beta isoform, beta isoform of regulatory subunit B56, protein phosphatase 2A, PP2A, B subunit, R5 beta isoform, protein phosphatase 2, regulatory subunit B\1, beta isoform, serine/threonine protein phosphatase 2A, 56 kDa regulatory subunit, beta isoform, PP2A, B subunit, PR61 beta isoform, PP2A, B subunit, B56 beta isoform, PP2A, B subunit, R5 beta isoform, PP2A, B subunit, B\1 beta isoform, protein phosphatase 2, regulatory subunit B (B56), beta isoform, PPP2R5B

Official Symbol: PPP2R5B

Accession Number(s): NP_006235.1

Human GeneID(s): [5526](#)

Immunogen

Peptide with sequence C-QRLTPQVAASGGQS, from the C Terminus of the protein sequence according to NP_006235.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx. 55kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 57.4kDa according to NP_006235.1). Recommended concentration: 2-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB05241 (2 μ g/ml) staining of Human Cerebellum lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.