



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07942 - Goat Anti-B7-H4 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** B7-H4, B7H4, B7S1, B7X, B7h.5, FLJ22418, PRO1291, RP11-229A19.4, VCTN1, T cell costimulatory molecule B7x, immune costimulatory protein, V-set domain containing T cell activation inhibitor 1

**Official Symbol:** VTCN1

**Accession Number(s):** NP\_078902.2; NP\_001240778.1

**Human GeneID(s):** [79679](#)

**Non-Human GeneID(s):** 242122 (mouse), 295322 (rat)

### Immunogen

Peptide with sequence C-SKGKGNANLEYK, from the internal region of the protein sequence according to NP\_078902.2; NP\_001240778.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:64000.

**Western blot:** Approx 26+21kDa bands observed in HEK293 cell lysates, which were both successfully blocked by incubation with the immunizing peptide. An approx 26kDa and 21kDa band observed in Human Adrenal Gland lysates, and approx 27kDa in Human Pancreas lysates (calculated MW of 30.9kDa according to NP\_078902.2 isoform 1 and 20.7kDa according to NP\_001240778.1 isoform 2 ). Recommended concentration: 0.1-1µg/ml. Primary incubation 1 hour at room temperature.


**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm and membranes/cell junctions of MCF7 cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of MCF7 cells. Recommended concentration: 10ug/ml.


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human, Mouse, Rat, Dog



EB07942 (1ug/ml) staining of HEK293 iso 1 + iso 2 cell lysate (A) + peptide (B). Human Adrenal Gland iso 1 (C), Human Adrenal Gland iso 2 (D) and (0.1µg/ml) Human pancreas iso 1 (E) lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB07942 Immunofluorescence analysis of paraformaldehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane/cell junction and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB07942 Flow cytometric analysis of paraformaldehyde fixed MCF7 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.