



Everest Biotech Ltd
Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD, United Kingdom

everestbiotech.com

sales@everestbiotech.com

support@everestbiotech.com

Tel +44 1869 238326

Fax +44 1869 238327

**Research Use Only. Not for
diagnostic or therapeutic use.**

Storage: For long-term storage
keep aliquots at -20°C. (Store no
longer than 12 months at 4°C).
Minimize freezing and thawing.

EB06344 - Goat Anti-BAF53A and BAF53B Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: hArpN beta, actin-related protein 4, INO80 complex subunit K, BRG1-associated factor, BAF53, BAF complex 53 kDa subunit, MGC5382, INO80K, Arp4, ARPN-BETA, 53 kDa BRG1-associated factor B, ACTL6B, ACTL6A, actin-related protein, hArpN alpha, actin-like 6B, actin-like 6A, BAF53b, BAF53a, ACTL6, BAF53B, BAF53A

Official Symbol: ACTL6A / ACTL6B

Accession Number(s): NP_004292.1; NP_817126.1; NP_057272.1

Human GeneID(s): [51412](#) , [86](#)

Important Comments: This antibody is expected to recognise both BAF53A and BAF53B, which are almost identical. Reported variants represent identical protein (NP_817126.1; NP_829888.1).

Immunogen

Peptide with sequence YEEGGKQCVERKCP, from the C Terminus of the protein sequence according to NP_004292.1; NP_817126.1; NP_057272.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 48kDa band observed in Hela lysates (predicted MW of 49kDa according to NP_057272, NP_004292). Recommended for use at 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow,



EB06344 staining (1 μ g/ml) of HeLa lysate (RIPA buffer, 35 μ g total protein per lane). Primary incubated for 1 hour.
Detected by western blot using chemiluminescence.