



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07132 - Goat Anti-BARK1 / GRK2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: BARK1, GRK2, ADRBK1, adrenergic, beta, receptor kinase 1, BETA-ARK1, beta adrenergic receptor kinase 1

Official Symbol: GRK2

Accession Number(s): NP_001610.2

Human GeneID(s): [156](#)

Non-Human GeneID(s): 110355 (mouse)

Immunogen

Peptide with sequence C-RDAYREAQQLVQR, from the internal region of the protein sequence according to NP_001610.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 80kDa band observed in Daudi cell lysates (calculated MW of 79.7kDa according to NP_001610.1). An additional band of unknown identity was observed in some lysates at 48kDa and was successfully blocked by incubation with the immunizing peptide. Recommended concentration: 0.3-2µg/ml. Primary incubation 1 hour at room temperature.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB07132 (2 μ g/ml) staining of Daudi cell lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.