

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08117 - Goat Anti-BDNF Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: BDNF, brain-derived neurotrophic factor, MGC34632, neurotrophin

Official Symbol: BDNF

Accession Number(s): NP_001700.2; NP_001700.2; NP_733928.1; NP_733929.1;

NP_733930.1; NP_733931.1 **Human GeneID(s):** 627

Important Comments: This antibody is expected to recognise all reported isoforms

 $(NP_001700.2; \, NP_001700.2; \, NP_733928.1; \, NP_733929.1; \, NP_733930.1; \, NP_733929.1; \, NP_733930.1; \, NP_733929.1; \, NP_$

NP_733931.1)

Immunogen

Peptide with sequence ETKCNPMGYTKE, from the internal region of the protein sequence according to NP_001700.2; NP_001700.2; NP_733928.1; NP_733929.1; NP_733930.1; NP_733931.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:2000.

Immunofluorescence: Expression of the protein seen in the nucleus and cytoplasm of Glioblastoma U251 cells. Recommended concentration: 10µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB08117 Immunofluorescence analysis of paraformaldehyde fixed U251 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing speckled nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).