



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05140-B - Goat Anti-CD20 / MS4A1 (C Terminus), biotinylated Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** MS4A1, membrane-spanning 4-domains, subfamily A, member 1, B1, Bp35, CD20, CVID5, LEU-16, MS4A2, S7, B-lymphocyte cell-surface antigen B1, CD20 antigen, CD20 receptor, leukocyte surface antigen Leu-16, MGC3969

**Official Symbol:** MS4A1

**Accession Number(s):** NP\_068769.2

**Human GeneID(s):** [931](#)

**Important Comments:** Reported variants represent identical protein: NP\_068769.2, NP\_690605.1

### Immunogen

Peptide with sequence CQDQESSPIENDSSP., from the C Terminus of the protein sequence according to NP\_068769.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:2000.

**Western blot:** Approx 35kDa band observed in lysates of cell line Daudi (calculated MW of 33.1kDa according to NP\_068769.2). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.3-1µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



Biotinylated EB05140 (0.3µg/ml) staining of Daudi lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.