



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12595 - Goat Anti-CD3-ZETA / CD247 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CD247, CD247 molecule, CD3-ZETA, CD3H, CD3Q, CD3Z, T3Z, TCRZ, CD247 antigen, zeta subunit, CD3Z antigen, zeta polypeptide (TIT3 complex), CD3zeta chain, T-cell antigen receptor complex, zeta subunit of CD3, T-cell receptor T3 zeta chain, T-cell surface glycoprotein CD3 zeta chain

Official Symbol: CD247

Accession Number(s): NP_932170.1; NP_000725.1

Human GeneID(s): [919](#)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_932170.1; NP_000725.1).

Immunogen

Peptide with sequence C-KNPQEGLYNELQKD, from the internal region of the protein sequence according to NP_932170.1; NP_000725.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 16-17kDa band observed in lysates of cell lines Jurkat and MOLT4 (calculated MW of 18.6kDa according to NP_000725.1). Recommended concentration: 0.1-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB12595 (0.1 μ g/ml) staining of Jurkat (A) and (0.3 μ g/ml) MOLT4 (B) cell lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.