

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB11932 - Goat Anti-CD57 / B3GAT1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** B3GAT1, beta-1,3-glucuronyltransferase 1 (glucuronosyltransferase P), CD57, GLCATP, GLCUATP, HNK1, LEU7, NK-1, NK1, LEU7 antigen, OTTHUMP00000235432, UDP-GlcUA:glycoprotein beta-1,3-glucuronyltransferase, galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1, glcUAT-P, glucuronosyltransferase P

**Official Symbol:** B3GAT1

**Accession Number(s):** NP\_061114.2

**Human GeneID(s):** [27087](#)

**Non-Human GeneID(s):** 76898 (mouse), 117108 (rat)

**Important Comments:** Reported variants represent identical protein: NP\_473366.1, NP\_061114.2.

### Immunogen

Peptide with sequence C-HVETPRNYKLRGD, from the internal region of the protein sequence according to NP\_061114.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:64000.

**Western blot:** Approx 38kDa band observed in Human Brain (Frontal Cortex) lysates (calculated MW of 38.3kDa according to NP\_061114.2). Recommended concentration: 0.1-0.3µg/ml. An additional band of 17kDa was consistently observed, however this band was not blocked by the immunizing peptide and it is therefore a non-specific signal. We call for caution when used for other assays than Western blot.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig



EB11932 (0.1 $\mu$ g/ml) staining of Human Frontal Cortex lysate (35 $\mu$ g protein in RIPA buffer) with (B) and without (A) blocking with the immunizing peptide. Primary incubation was 1 hour. Detected by chemiluminescence.