



Everest Biotech Ltd
Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD, United Kingdom

everestbiotech.com

sales@everestbiotech.com

support@everestbiotech.com

Tel +44 1869 238326

Fax +44 1869 238327

Research Use Only. Not for diagnostic or therapeutic use.

Storage: For long-term storage keep aliquots at -20°C. (Store no longer than 12 months at 4°C). Minimize freezing and thawing.

EB06123 - Goat Anti-CENTB2 / ACAP2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ArfGAP with coiled-coil, ankyrin repeat and PH domains 2, Arf GAP with coiled coil, ANK repeat and PH domains 2, centaurin beta2, KIAA0041, CNT-B2, centaurin, beta 2, CENTB2, ACAP2

Official Symbol: ACAP2

Accession Number(s): NP_036419.3

Human GeneID(s): [23527](#)

Non-Human GeneID(s): 78618 (mouse)

Immunogen

Peptide with sequence C-EKLNRFQQDSQKF, from the C Terminus of the protein sequence according to NP_036419.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Approx 90kDa band observed in Human T cell line MOLT4 lysates (calculated MW of 88.1kDa according to NP_036419). Recommended concentration: 1-3µg/ml.

IHC: In paraffin embedded Human Lung shows strong staining of Type II pneumocytes. Recommended concentration, 5-10µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig



EB06123 (1 μ g/ml) staining of MOLT4 lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.



EB06123 (5 μ g/ml) staining of paraffin embedded Human Lung. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.