

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07350 - Goat Anti-CILP Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CILP, cartilage intermediate layer protein, nucleotide

pyrophosphohydrolase, HGNC:1980, HsT18872, cartilage intermediate layer protein

Official Symbol: CILP

Accession Number(s): NP_003604.2

Human GenelD(s): 8483

Non-Human GenelD(s): 214425 (mouse), 315761 (rat)

Immunogen

Peptide with sequence C-NYRRTDHEDPRVK, from the internal region of the protein sequence according to NP_003604.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 150kDa band observed in rat spinal cord lysates (calculated MW of

133kDa according to human NP_003604.2 and of 132kDa according to Rat

NP_001101631.1). Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB07350 ($0.03\mu g/ml$) staining of rat spinal cord lysate ($35\mu g$ protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.