



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB10236-B - Goat Anti-CIRBP (aa 161-172), Biotinylated Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CIRBP, cold inducible RNA binding protein, CIRP, A18 hnRNP, cold inducible RNA-binding protein, glycine-rich RNA binding protein, Cold-inducible RNA-binding protein

**Official Symbol:** CIRBP

**Accession Number(s):** NP\_001271.1

**Human GeneID(s):** [1153](#)

**Non-Human GeneID(s):** 12696 (mouse), 81825 (rat)

**Important Comments:** This antibody is expected to recognize reported isoform 1 (NP\_001271.1) only.

### Immunogen

Peptide with sequence C-RDSYDSYATHNE., from the C Terminus of the protein sequence according to NP\_001271.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 18kDa band observed in Rat Testis lysates (calculated MW of 18.6kDa according to Rat NP\_112409.2). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 1-3µg/ml.

### Species Reactivity

**Tested:** Human, Mouse, Rat

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow



Biotinylated EB10236 (1 $\mu$ g/ml) staining of Rat Testis lysate (35 $\mu$ g protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.