

### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08544 - Goat Anti-CLEC2 / CLEC1B Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** CLEC1B, CLEC2, C-type lectin domain family 1, member B, 1810061I13Rik, CLEC2B, PRO1384, QDED721, C-type lectin-like receptor-2

Official Symbol: CLEC1B

Accession Number(s): NP\_001092901.1; NP\_057593.3

Human GeneID(s): 51266

Important Comments: This antibody is expected to recognize both reported isoforms

(NP\_001092901.1; NP\_057593.3)

## **Immunogen**

Peptide with sequence KGHKCSPCDTNWRY, from the internal region of the protein sequence according to NP\_001092901.1; NP\_057593.3.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:1000.

**Western blot:** Approx 28kDa band observed in human bone marrow and human kidney lysates (calculated MW of 26.6kDa according to NP\_001092901.1). Recommended

concentration: 1-3µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB08544 ( $1\mu g/ml$ ) staining of human kidney lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.