

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB12398 - Goat Anti-CLIC1 / NCC27 (aa45-57) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CLIC1, chloride intracellular channel 1, G6, NCC27, RNCC protein, chloride channel ABP, chloride intracellular channel protein 1, hRNCC, nuclear chloride ion channel 27, nuclear chloride ion channel protein, p64CLCP, regulatory nuclear chloride ion channe

Official Symbol: CLIC1

Accession Number(s): NP_001279.2

Human GeneID(s): 1192

Non-Human GenelD(s): 114584 (mouse), 406864 (rat)

Immunogen

Peptide with sequence C-TVDTKRRTETVQK, from the internal region of the protein sequence according to NP_001279.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.

Western blot: Approx 30kDa band observed in lysates of cell line U937 (calculated MW of 26.9kDa according to NP_001279.2). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB12398 (2μg/ml) staining of U937 lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour.

Detected by chemiluminescence.