

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB12399 - Goat Anti-CLIC1 / NCC27 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CLIC1, chloride intracellular channel 1, G6, NCC27, RNCC protein, chloride channel ABP, chloride intracellular channel protein 1, hRNCC, nuclear chloride ion channel 27, nuclear chloride ion channel protein, p64CLCP, regulatory nuclear chloride ion channe Official Symbol: CLIC1 Accession Number(s): NP_001279.2 Human GeneID(s): <u>1192</u> Non-Human GeneID(s): 114584 (mouse), 406864 (rat)

Immunogen

Peptide with sequence C-TEVHTDTNKIEE, from the internal region of the protein sequence according to NP_001279.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 30kDa band observed in lysates of cell line U937 (calculated MW of 26.9kDa according to NP_001279.2). Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Colon. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB12398 (2µg/ml) staining of U937 lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB12399 (5µg/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.