



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08147 - Goat Anti-COL4A3BP (aa662-672) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: COL4A3BP, collagen, type IV, alpha 3 (Goodpasture antigen) binding protein, CERT, CERTL, FLJ20597, GPBP, STARD11, Goodpasture antigen-binding protein, START domain containing 11, alpha 3 type IV collagen binding protein, ceramide transporter

Official Symbol: CERT1

Accession Number(s): NP_005704.1; NP_112729.1

Human GeneID(s): [10087](#)

Non-Human GeneID(s): 68018 (mouse)

Important Comments: This antibody is expected to recognise both reported isoforms (NP_005704.1 and NP_112729.1)

Immunogen

Peptide with sequence DHDSAPLNNRC, from the internal region of the protein sequence according to NP_005704.1; NP_112729.1 .

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 60-65kDa band observed in Mouse Heart lysates (calculated MW of 71.1kDa according to Mouse NP_075909.1). An additional band of unknown identity was also consistently observed at 25kDa. Both bands were successfully blocked by incubation with the immunising peptide. Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.

Species Reactivity

Tested: Mouse

Expected from sequence similarity: Human, Mouse



EB08147 (0.5µg/ml) staining of Mouse Heart lysate (A) + peptide (B) (35µg protein in RIPA buffer). Detected by chemiluminescence.