

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06781 - Goat Anti-Collagen, type VII Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: Collagen, type VII, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive), COL7A1, HGNC:2214, EBD1, EBDCT, EBR1, LC collagen, alpha 1 type VII collagen, collagen VII, alpha-1 polypeptide, long chain collagen, collagen, type VII, alpha 1 Official Symbol: COL7A1 Accession Number(s): NP_000085.1 Human GeneID(s): 1294 Non-Human GeneID(s): 12836 (mouse) Important Comments: This product does not work on cryostat sections of Normal Human Skin.

Immunogen

Peptide with sequence C-TVQYSDDPRTEF, from the internal region of the protein sequence according to NP_000085.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000. **Western blot:** Approx 250kDa band observed in Rat Testis lysates (calculated MW of 159kDa according to Rat NP_001100328.1). Recommended concentration: 0.1-1.5µg/ml.

A customer found this antibody unsuccessful on cryosections of Human Skin.

Species Reactivity

Tested: Rat Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig

250kDa 150kDa 100kDa 75kDa
50kDa
37kDa
25kDa
20kDa

15kDa

EB06781 (0.5µg/ml) staining of Rat Testis lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.