



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09614 - Goat Anti-CYB5R3 / Dia 1 (mouse) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** Cyb5r3, cytochrome b5 reductase 3, 0610016L08Rik, 2500002N19Rik, C85115, Dia-1, Dia1, WU:AL591952.1-001, WU:AL591952.1-002, WU:AL591952.1-003, WU:Cyb5r3, NADH-cytochrome b5 reductase, diaphorase 1, diaphorase 1 (NADH)

**Official Symbol:** Cyb5r3

**Accession Number(s):** NP\_084063.1

**Non-Human GeneID(s):** 109754 (mouse), 25035 (rat)

### Immunogen

Peptide with sequence PNLERVGHPKERC, from the C Terminus of the protein sequence according to NP\_084063.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.


**Western blot:** Approx 35kDa band observed in Mouse Liver lysates as well as in mouse and rat brain lysates (calculated MW of 34.1kDa according to mouse NP\_084063.1).

Recommended concentration: 0.1-1µg/ml.

### Species Reactivity

**Tested:** Mouse

**Expected from sequence similarity:** Mouse, Rat, Dog, Cow



EB09614 (0.1 $\mu$ g/ml) staining of Mouse Liver lysate (RIPA buffer, 35 $\mu$ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.