

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: <u>sales@everestbiotech.com</u> Tech support: <u>support@everestbiotech.com</u>

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10306 - Goat Anti-CYPOR Antibody

Size: 100µg specific antibody in 200µl

C

Target Protein

Principal Names: CPR, CYPOR, cytochrome P450 reductase, DKFZp686G04235, FLJ26468, NADPH-dependent cytochrome P450 reductase, P450 (cytochrome) oxidoreductase, P450R, POR Official Symbol: POR Accession Number(s): NP_000932.3 Human GenelD(s): <u>5447</u> Non-Human GenelD(s): 18984 (mouse), 29441 (rat)

Immunogen

Peptide with sequence C-TNPPRTNVLYE, from the internal region of the protein sequence according to NP_000932.3.

Please note the peptide is available for sale.

Purification and Storage


Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:16000.
Western blot: Approx 90kDa band observed in Human, Mouse and Rat Liver lysates (calculated MW of 77.0kDa according to NP_000932.3). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.
IHC: Paraffin embedded Human Adrenal Gland. Recommended concentration: 3.75µg/ml.

Species Reactivity

Tested: Human, Mouse, Rat **Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow, Pig



EB10306 (0.1µg/ml) staining of Rat Liver lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB10306 (3.75µg/ml) staining of paraffin embedded Human Adrenal Gland. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.