



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11100 - Goat Anti-D-amino-acid oxidase (aa286-298) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: DAAO, D-amino-acid oxidase, DAMOX, MGC35381, OXDA, DAO

Official Symbol: DAO

Accession Number(s): NP_001908.3

Human GeneID(s): [1610](#)

Immunogen

Peptide with sequence C-RPQIRLEREQLRT, from the internal region of the protein sequence according to NP_001908.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 37kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 39.5kDa according to NP_001908.3). Recommended concentration: 0.03-0.01µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog, Pig, Cow

Specific References

This antibody has been successfully used in WB on Macaques:

Nuzzo T, Punzo D, Devoto P, Rosini E, Paciotti S, Sacchi S, Li Q, Thiolat ML, Véga C, Carella M, Carta M, Gardoni F, Calabresi P, Pollegioni L, Bezard E, Parnetti L, Errico F, Uziel A

The levels of the NMDA receptor co-agonist D-serine are reduced in the substantia nigra of MPTP-lesioned macaques and in the cerebrospinal fluid of Parkinson's disease patients
Sci Rep. 2019 Jun 20;9(1):8898.

PMID: 31222058


This antibody has been successfully used in WB and IF on Mouse:

Sasabe J, Miyoshi Y, Rakoff-Nahoum S, Zhang T, Mita M, Davis BM, Hamase K, Waldor MK

Interplay between microbial D-amino acids and host D-amino acid oxidase modifies murine mucosal defence and gut microbiota

Nat Microbiol. 2016 Jul 25;1(10):16125

PMID: 27670111



EB11100 (0.03 μ g/ml) staining of Human Cerebellum lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.