

### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10930 - Goat Anti-Dbf4 (mouse) Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: Dbf4, DBF4 homolog (S. cerevisiae), AA545217, Ask, activator of S

phase kinase, protein DBF4 homolog A

Official Symbol: Dbf4

Accession Number(s): NP\_038754.1; NP\_001177646.1 Non-Human GenelD(s): 27214 (mouse), 312046 (rat)

Important Comments: This antibody is expected to recognize both reported isoforms

(NP\_038754.1; NP\_001177646.1).

### **Immunogen**

Peptide with sequence C-DKKRTEYLPAHEDRT, from the internal region of the protein sequence according to NP\_038754.1; NP\_001177646.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

**Western blot:** Approx 70kDa band observed in Mouse and Rat Testis and Thymus lysates (calculated MW of 74.2kDa according to NP\_038754.1). Recommended

concentration: 1-3µg/ml.

### **Species Reactivity**

Tested: Mouse, Rat

Expected from sequence similarity: Mouse, Rat

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB10930 ( $1\mu g/ml$ ) staining of Mouse Testis lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.