



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12413 - Goat Anti-DDX41 (aa468-480) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: DDX41, DEAD (Asp-Glu-Ala-Asp) box polypeptide 41, ABS, 2900024F02Rik, DEAD box protein 41, DEAD box protein abstract homolog, DEAD-box protein abstract, probable ATP-dependent RNA helicase DDX41, putative RNA helicase

Official Symbol: DDX41

Accession Number(s): NP_057306.2

Human GeneID(s): [51428](#)

Non-Human GeneID(s): 72935 (mouse), 314336 (rat)

Immunogen

Peptide with sequence C-QEERTKAIEAFRE, from the internal region of the protein sequence according to NP_057306.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.


Western blot: Approx 85kDa band observed in Human Bone Marrow lysates (calculated MW of 69.8kDa according to NP_057306.2). This molecular weight is routinely observed by other sources. Recommended concentration: 0.5-2µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB12413 (0.5µg/ml) staining of Human Bone Marrow lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB12413 (5µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.