



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09819 - Goat Anti-DYDC1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: bA36D19.5, BC019250 protein, DPY30 domain containing 1, DPY30D1, FLJ43920, OTTHUMP00000019963, DYDC1

Official Symbol: DYDC1

Accession Number(s): NP_620167.1

Human GeneID(s): [143241](#)

Immunogen

Peptide with sequence C-RNEDILHSEEATLD, from the internal region of the protein sequence according to NP_620167.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: In transfected HEK293 transiently expressing Human DYDC1 (myc and DYKDDDDK tagged), a band of approx. 28kDa is observed. No bands are observed in mock-transfected HEK293 and the same band is observed using anti-myc antibody. Recommended concentration, 0.5-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



Lane A - EB09819 (1 μ g/ml) staining of HEK293 overexpressing Human DYDC1 lysate (10 μ g protein in RIPA buffer) Lane B - EB09819 (1 μ g/ml) staining of HEK293 mock-transfected lysate (10 μ g protein in RIPA buffer). Lane C - EB00106 anti-MYC Tag (1/1000) staining HEK293 overexpressing Human DYDC1 lysate (10 μ g protein in RIPA buffer). Primary incubations were for 1 hour. Detected by chemiluminescence.