



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB13023 - Goat Anti-FABP1 / L-FABP (aa100-112) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** FABP1, fatty acid binding protein 1, liver, FABPL, L-FABP, fatty acid-binding protein 1, fatty acid-binding protein, liver, liver-type fatty acid-binding protein

**Official Symbol:** FABP1

**Accession Number(s):** NP\_001434.1

**Human GeneID(s):** [2168](#)

### Immunogen

Peptide with sequence C-SVTELNGDIITNT, from the internal region of the protein sequence according to NP\_001434.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:128000.

**Western blot:** Approx 14kDa band observed in Human and Pig Liver lysates (calculated MW of 14.2kDa according to Human NP\_001434.1 and 14.1kDa according to Pig NP\_001004046.1). Recommended concentration: 0.03-0.1µg/ml.

### Species Reactivity

**Tested:** Human, Pig

**Expected from sequence similarity:** Human, Dog, Pig



EB13023 (0.1 $\mu$ g/ml) staining of Human (A) and Pig (B) Liver lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.