



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12000 - Goat Anti-FABP3 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: FABP3, fatty acid binding protein 3, muscle and heart (mammary-derived growth inhibitor), FABP11, H-FABP, M-FABP, MDGI, O-FABP, Fatty acid-binding protein 3, muscle, fatty acid binding protein 11, fatty acid-binding protein, heart, heart-type fatty acid-binding protein, mammary-derived growth inhibitor, muscle fatty acid-binding protein

Official Symbol: FABP3

Accession Number(s): NP_004093.1

Human GeneID(s): [2170](#)

Immunogen

Peptide with sequence C-PTTIEKNGDILTLK, from the internal region of the protein sequence according to NP_004093.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 15kDa band observed in Human, Mouse and Rat Heart lysates (calculated MW of 14.9kDa according to NP_004093.1). Recommended concentration: 0.01-0.03µg/ml.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Cow



EB12000 (0.01 μ g/ml) staining of Human (A), Mouse (B) and Rat(C) Heart lysates (35 μ g protein in RIPA buffer).
Primary incubation was 1 hour. Detected by chemiluminescence.