

## **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB12004 - Goat Anti-FABP4 Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** FABP4, fatty acid binding protein 4, adipocyte, A-FABP, AFABP, ALBP, aP2, adipocyte lipid-binding protein, adipocyte-type fatty acid-binding protein, fatty

acid-binding protein 4, fatty acid-binding protein, adipocyte **Official Symbol:** FABP4

Accession Number(s): NP\_001433.1

Human GeneID(s): 2167

### **Immunogen**

Peptide with sequence C-SVNGDVITIKSEST, from the internal region of the protein sequence according to NP\_001433.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 14kDa band observed in Human Adipose tissue lysates (calculated MW of 14.7kDa according to NP\_001433.1). Recommended concentration: 0.03-0.1µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Dog

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

EB12004 (0.03μg/ml) staining of Human Adipose lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.