



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07823 - Goat Anti-G6PD (aa 305 - 318) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: G6PD, glucose-6-phosphate dehydrogenase, G6PD1, glucose-6-phosphate 1-dehydrogenase, glucose-6-phosphate dehydrogenase, G6PD

Official Symbol: G6PD

Accession Number(s): NP_000393.4 ; NP_001035810.1

Human GeneID(s): [2539](#)

Non-Human GeneID(s): 14381 (mouse), 24377 (rat)

Important Comments: This antibody is expected to recognise both reported isoforms (NP_000393.4 and NP_001035810.1).

Immunogen

Peptide with sequence C-KPASTNSDDVRDEK, from the internal region of the protein sequence according to NP_000393.4 ; NP_001035810.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 55kDa band observed in Human Peripheral Blood Mononucleocyte lysates (calculated MW of 59.3kDa according to NP_001035810.1). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07823 (1µg/ml) staining of Peripheral Blood Mononucleocytes lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.