

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08257 - Goat Anti-C2GnT-M (aa 273 to 284) Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: GCNT3, glucosaminyl (N-acetyl) transferase 3, mucin type, C2/4GnT,

C2GnT-M, C2GnT2, GnT-M, beta1,6-N-acetylglucosaminyltransferase

Official Symbol: GCNT3

Accession Number(s): NP\_004742.1

Human GeneID(s): 9245

### **Immunogen**

Peptide with sequence C-RDTLHLTNKKKD, from the internal region of the protein sequence according to NP\_004742.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:2000.

**Western blot:** Approx. 50kDa band observed in Human Lymph Node lysates (calculated MW of 50.9kDa according to NP\_004742.1). In transfected HEK293 transiently expressing GCNT3 a band of approx. 55kDa is observed. This band is not observed in the non-transfected HEK293. Recommended concentration: 0.3-1µg/ml.


## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB08257 (0.3 $\mu$ g/ml) staining of human lymph node lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



HEK293 overexpressing GCNT3 (RC202007) and probed with EB08257 (mock transfection in first lane), tested by Origene.