

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB09243 - Goat Anti-GRM7 Antibody

Size: 100µg specific antibody in 200µl



Target Protein Principal Names: GRM7, glutamate receptor, metabotropic 7, FLJ40498, GLUR7, GPRC1G, MGLUR7, mGlu7 Official Symbol: GRM7 Accession Number(s): NP_000835.1; NP_870989.1 Human GenelD(s): 2917 Non-Human GenelD(s): 108073 (mouse), 81672 (rat) Important Comments: This antibody is expected to recognize both reported isoforms (NP_000835.1; NP_870989.1).

Immunogen

Peptide with sequence NCKLTISGSKKEDT, from the internal region of the protein sequence according to NP_000835.1; NP_870989.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000. **Western blot:** Approx. 100kDa band observed in Human Brain (Cerebellum) and in Mouse and Rat Brain lysates (calculated MW of 102kDa according to NP_000835.1). Recommended concentration 0.5-2µg/ml.

Species Reactivity

Tested: Human Expected from sequence similarity: Human, Mouse, Rat

-	250kDa 150kDa 100kDa 75kDa
	50kDa
	37kDa
	25kDa
	20kDa
	15kDa

EB09243 (0.5µg/ml) staining of Human Cerebellum lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.