

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10175 - Goat Anti-HCN3 (aa 715-728) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: hyperpolarization activated cyclic nucleotide-gated potassium channel 3, KIAA1535, MGC131493, potassium/sodium hyperpolarization-activated cyclic

nucleotide-gated channel 3, HCN3

Official Symbol: HCN3

Accession Number(s): NP_065948.1

Human GenelD(s): 57657

Non-Human GenelD(s): 15168 (mouse), 114245 (rat)

Immunogen

Peptide with sequence C-SQPSLPQRATGDGS, from the internal region of the protein sequence according to NP_065948.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 80kDa band observed in Mouse Brain lysates (calculated MW of 86.6kDa according to Mouse NP_032253.1). Recommended concentration: 0.01-0.03µg/ml.

Species Reactivity

Tested: Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow, Pig

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

15kDa

EB10175 ($0.01\mu g/ml$) staining of Mouse Brain lysate ($35\mu g$ protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.