



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09957 - Goat Anti-Herculin / MYF6 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: bHLHc4, herculin, MRF4, myogenic factor 6, myogenic factor 6 (herculin), myogenic factor-6, MYF6

Official Symbol: MYF6

Accession Number(s): NP_002460.1

Human GeneID(s): [4618](#)

Non-Human GeneID(s): 17878 (mouse), 25714 (rat)

Immunogen

Peptide with sequence C-SYRPKQENLEGAD, from the internal region of the protein sequence according to NP_002460.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: In transfected HEK293 transiently expressing full-length Human MYF6 (myc and DYKDDDDK tagged), a band of approx. 37kDa was observed. No bands were observed in mock-transfected HEK293 and the same band was observed using anti-myc tag antibody. Recommended concentration, 0.5-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Cow, Pig



HEK293 lysate (10ug protein in RIPA buffer) overexpressing Human MYF6 with MYC tag probed with EB09957 (0.5ug/ml) in Lane A and probed with anti-MYC Tag (1/1000) in lane C. Mock-transfected HEK293 probed with EB09957 (0.5mg/ml) in Lane B. Primary incubations were for 1 hour. Detected by chemiluminescence.