



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09623 - Goat Anti-HMGI-C / HMGA2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: HMGA2, high mobility group AT-hook 2, BABL, HMGI-C, HMGIC, LIPO, STQTL9, High-mobility group protein HMGI-C, high-mobility group (nonhistone chromosomal) protein isoform I-C

Official Symbol: HMGA2

Accession Number(s): NP_003474.1; NP_003475.1

Human GeneID(s): [8091](#)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_003474.1; NP_003475.1) and is not expected to cross-react with HMGA1.

Immunogen

Peptide with sequence C-KAAQKKAETGEK, from the internal region of the protein sequence according to NP_003474.1; NP_003475.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 18kDa band observed in Human Heart lysates (calculated MW of 11.8kDa according to NP_003474.1 and 11.5kDa according to NP_003475.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Di Cello et al, Mol Cancer Res. 2008 May;6(5):743-50.; PMID: 18505920). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog



EB09623 (1µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.