



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12832 - Goat Anti-HSD17B3 (aa141-155) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: HSD17B3, hydroxysteroid (17-beta) dehydrogenase 3, EDH17B3, SDR12C2, 17-beta-HSD 3, 17-beta-HSD3, 17-beta-hydroxysteroid dehydrogenase type 3, short chain dehydrogenase/reductase family 12C, member 2, testicular 17-beta-hydroxysteroid dehydrogenase, testo

Official Symbol: HSD17B3

Accession Number(s): NP_000188.1

Human GeneID(s): [3293](#)

Immunogen

Peptide with sequence C-SHFLNAPDEIQSLIH, from the internal region of the protein sequence according to NP_000188.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 33kDa band observed in Human Testis lysates (calculated MW of 34.5kDa according to NP_000188.1). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB12832 (2 μ g/ml) staining of Human Testis lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.