



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12522 - Goat Anti-IL3RA / CD123 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: IL3RA, interleukin 3 receptor, alpha (low affinity), CD123, IL3R, IL3RAY, IL3RX, IL3RY, hIL-3Ra, CD123 antigen, IL-3 receptor alpha SP2 isoform, IL-3 receptor subunit alpha, IL-3R subunit alpha, IL-3R-alpha, IL-3RA, interleukin-3 receptor subunit alpha

Official Symbol: IL3RA

Accession Number(s): NP_002174.1; NP_001254642.1

Human GeneID(s): [3563](#)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_002174.1; NP_001254642.1).

Immunogen

Peptide with sequence RQQYECLHYKTD, from the internal region of the protein sequence according to NP_002174.1; NP_001254642.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.


Western blot: Approx 43kDa band observed in lysates of cell lines Jurkat and Molt4 (calculated MW of 43.3kDa according to NP_002174.1). Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Colon. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB12522 (2 μ g/ml) staining of Jurkat (A) and Molt4 (B) lysates (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB12522 (5 μ g/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.