



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09548 - Goat Anti-INDOL1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: IDO2, indoleamine 2,3-dioxygenase 2, INDOL1, indoleamine 2,3-dioxygenase-like 1 protein, indoleamine-pyrrole 2,3 dioxygenase-like 1

Official Symbol: IDO2

Accession Number(s): NP_919270.3; NP_001382135.1

Human GeneID(s): [169355](#)

Immunogen

Peptide with sequence C-RDKTLESILHPR, from the C Terminus of the protein sequence according to NP_919270.3; NP_001382135.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx. 50kDa band observed in Human Kidney, Liver, Placenta and Tonsil lysates (calculated MW of 47.07kDa according to NP_919270.3). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog, Cow



EB09548 optimised QC. Primary incubation 1 hour at room temperature.
 Image A: Human Kidney lysate at primary Ab concentration 2µg/ml, Images B, C, and D: Human Placenta, Tonsil and Liver lysate at primary Ab concentration 3µg/ml. (Loaded 35µg protein in RIPA buffer, per lane). Detected by chemiluminescence.