



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08068 - Goat Anti-KCNQ4 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: KCNQ4, potassium voltage-gated channel, KQT-like subfamily, member 4, DFNA2, KV7.4, potassium channel KQT-like 4, potassium voltage-gated channel KQT-like protein 4

Official Symbol: KCNQ4

Accession Number(s): NP_004691.2; NP_751895.1

Human GeneID(s): [9132](#)

Important Comments: This antibody is expected to recognise both reported isoforms (NP_004691.2; NP_751895.1); may cross-react in Mouse;

Immunogen

Peptide with sequence C-DKGPSDAEVVDE, from the internal region of the protein sequence according to NP_004691.2; NP_751895.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.

Western blot: Approx 70kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 71.2kDa according to NP_751895.1). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB08068 (2µg/ml) staining of Cerebellum lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.