



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10963 - Goat Anti-KEAP1 (aa41-53) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: KEAP1, kelch-like ECH-associated protein 1, INrf2, KIAA0132, KLHL19, MGC10630, MGC1114, MGC20887, MGC4407, cytosolic inhibitor of Nrf2, kelch-like protein 19

Official Symbol: KEAP1

Accession Number(s): NP_036421.2

Human GeneID(s): [9817](#)

Non-Human GeneID(s): 50868 (mouse), 117519 (rat)

Important Comments: Reported variants represent identical protein: NP_036421.2, NP_987096.1

Immunogen

Peptide with sequence C-EVTPSQHGNRTFS, from the internal region (near N Terminus) of the protein sequence according to NP_036421.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


IHC: In paraffin embedded Human Cerebellum shows textured cytoplasm staining in Purkinje cell and processes. Recommended concentration: 3-5µg/ml.

Immunofluorescence: Strong expression of the protein seen in the nucleus and cytoplasm of HeLa cells. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB10963 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (5ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (5ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB10963 (3.8µg/ml) staining of paraffin embedded Human Cerebellum. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.