

#### **UK Office**

**Everest Biotech Ltd** 

Cherwell Innovation Centre

77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10533 - Goat Anti-KLHDC8B Antibody

Size: 100µg specific antibody in 200µl



**Target Protein** 

Principal Names: FLJ11302, kelch domain containing 8B, kelch domain-containing

protein 8B, MGC35097, KLHDC8B **Official Symbol:** KLHDC8B

Accession Number(s): NP\_775817.1

Human GeneID(s): 200942

Non-Human GenelD(s): 78267 (mouse), 306589 (rat)

#### **Immunogen**

Peptide with sequence C-TKLPRSLRMRDK, from the internal region of the protein sequence according to NP\_775817.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 38kDa band observed in lysates of cell lines Daudi, Jurkat and NIH3T3 (calculated MW of 37.7kDa according to NP\_775817.1). Recommended concentration: 0.5-2µg/ml. An additional band of unknown identity was also consistently observed at 22kDa. This band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

#### **Species Reactivity**

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB10533 ( $0.5\mu g/ml$ ) staining of Daudi lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.