

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09262 - Goat Anti-LAT1 / SLC7A5 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SLC7A5, solute carrier family 7 (cationic amino acid transporter, y+ system), member 5, 4F2LC, CD98, D16S469E, E16, LAT1, MPE16, hLAT1, 4F2 light chain, CD98 light chain, L-type amino acid transporter 1, integral membrane protein E16, large neutral amino acids transporter 1, large neutral amino acids transporter small subunit 1, sodium-independent neutral amino acid transporter LAT1

Official Symbol: SLC7A5

Accession Number(s): NP_003477.4

Human GeneID(s): [8140](#)

Immunogen

Peptide with sequence C-RHRKPELERPIKVN, from the internal region of the protein sequence according to NP_003477.4.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 60kDa band observed in Human Spleen lysates (calculated MW of 55.0kDa according to NP_003477.4). Recommended concentration: 1-3µg/ml. Some minor background is detected and is blocked by the immunizing peptide. We call for caution when used for other assays than Western blot.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog



EB09262 (0.3µg/ml) staining of Human Spleen lysate (35µg protein in RIPA buffer) with (B) and without (A) blocking with the immunising peptide. Primary incubation was 1 hour. Detected by chemiluminescence.