

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12997 - Goat Anti-LIMK2 (aa181-193) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: LIMK2, LIM domain kinase 2

Official Symbol: LIMK2

Accession Number(s): NP_001026971.1; NP_005560.1; NP_057952.1

Human GeneID(s): [3985](#)

Non-Human GeneID(s): 16886 (mouse), 29524 (rat)

Important Comments: This antibody is expected to recognize all reported isoforms (NP_001026971.1; NP_005560.1; NP_057952.1).

Immunogen

Peptide with sequence C-QVKEVNRMHISPN, from the internal region of the protein sequence according to NP_001026971.1; NP_005560.1; NP_057952.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 75+80kDa doublet band observed in lysates of cell lines HeLa and NIH3T3 and approx 70kDa observed in Rat Pancreas lysates (calculated MW of 77.9kDa according to Human NP_001026971.1 and 72.2kDa according to Human NP_005560.1, Mouse NP_034848.1 and Rat NP_077049.2). Recommended concentration:0.3-1µg/ml.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Pig



EB12997 (1 μ g/ml) staining of HeLa (A) and NIH3T3 (B) lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB12997 (0.3 μ g/ml) staining of Rat Pancreas lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.