



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11790 - Goat Anti-MCTS1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: FLJ39637, malignant T cell amplified sequence 1, malignant T cell-amplified sequence 1, MCT1, MCT-1, MCTS1, multiple copies T-cell malignancies, OTTHUMP00000023950, OTTHUMP00000023951

Official Symbol: MCTS1

Accession Number(s): NP_054779.1; NP_001131026.1

Human GeneID(s): [28985](#)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_054779.1; NP_001131026.1).

Immunogen

Peptide with sequence DEKENVSNCIQLKTS, from the N Terminus of the protein sequence according to NP_054779.1; NP_001131026.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 19kDa band observed in lysates of cell lines Jurkat and MOLT4 (calculated MW of 20.6kDa according to NP_054779.1 and NP_001131026.1).

Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog, Pig, Cow



EB11790 (0.3µg/ml) staining of MOLT4 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.